

Borehole Information

Borehole Number:

105/25/K003/1A/H4

Prospecting Company: De Beers Holdings Botswana

Prospecting License No.: PL31/88

Coordinates Type: WGS 84

Coordinates Format: Decimal Degrees

Coordinates: -22.704422 , 27.883914

Location: Lerala

Start Drill: N/A

End Drill: N/A

Borehole Type: Exploration

Drilling Method: Core

Sample Type: Core

Depth of Surface Casing: N/A

Azimuth: Degrees

Hole Inclination: N/A

Collar Elevation: N/A

Borehole Water Strike: N/A

End of Hole (Depth): N/A

Log Formation:

No.	From (m)	To (m)	Log Formation
1.	0	1	Brown Topsoil.
2.	1	7	Clay fraction is yellowish in colour. Weak calcretisation slightly silicified, highly weathered light brown coarsely textured kimberlite with inclusions of Amphibolitebolite, quartz and k-feldspar 2m - overall brown/red staining.
3.	7	11	Clay fraction is greenish in colour, 8m - 99% inclusions (qtz & pink kfs coated is greenish/yellow material) 10m-20% inclusions.
4.	11	18	20% inclusions.
5.	18	19	Apple/olive, phlogopite bearing, olive poor coarsely textured, generally featureless hypabyssal kimberlite. Inclusions rich - variable proportions.
6.	19	20	95% inclusions (sheared greenish chloritic qtz, feldspar, kfs)
7.	20	21	100% inclusions (chloritised QF gneiss).
8.	21	29	80% inclusions (pink kfs)
9.	29	30	40% inclusions
10.	30	39	30% inclusions (kfs, Amphibolitebolite).
11.	39	44	Below water table kimberlite is fresher, deeper olive green. Coarsely textured, featureless, phlogopite bearing, olivine poor, hypabyssal kimberlite macrocrystic phlog up to 3mm observed. 40% inclusions (kfs Amphibolitebolite)
12.	44	47	40% inclusions (chlorite qtz, kfs, Amphibolitebolite),


Physical Address: Plot 11566, Khama 1 Avenue, Lobatse, Botswana

Postal Address: Private Bag 0014, Lobatse, Botswana

Telephone: +267 533 0327 / +267 533 0428 **Fax:** +267 533 4295

Email: info@bgi.org.bw **Website:** www.bgi.org.bw

Page Number: 1 | **Date/Time Stamp:** Fri, 18 Oct 24 06:11:25 +0200





Borehole Information

13.	47	48	30% inclusions (chlorite qtz, kfs, Amphiboliteib).
14.	48	49	70% inclusions.
15.	49	54	49m notable chloritisation of gneisses inclusions.
16.	54	59	90% inclusions (chloritic sheared qtz bearing rock + kfs).
17.	59	65	20% inclusions (chlorite qtz, kfs, Amphiboliteibolite),
18.	65	66	10% inclusions.
19.	66	67	5% inclusions.
20.	67	69	65m - distinct change to coloured inclusions fresh kimberlite. Olive green/grey featureless, coarsely textured hypabyssal kimberlite.
21.	69	71	5% inclusions 70m.
22.	71	77	Basically inclusion fresh down to 90m.
23.	77	89	Distinct lack of the usual coloured inclusions qtz, kfs.
24.	89	90	Appears to be 100% grey, features hypabyssal kimberlite at first glance, but there is a high proportion of fine grained biotite Schist. Visually confusing and difficult to separate from kimberlite.
25.	90	94	94m predominantly Amphiboliteibolite 1x2mm phlog obs.
26.	94	95	50% inclusions (Amphiboliteibolite).
27.	95	96	Mostly biotite Schist is some Amphiboliteibolite and pink gneiss. Phlogopite observed in fine drill product (could be bt). Visually confusing difficult to separate kimberlite.

NB: For any enquiries, application, acquisition or verifications related to Water Boreholes, kindly contact the Department of Water and Sanitation for assistance.

